=> file .meeting

'EVENTLINE' IS NOT A VALID FILE NAME

Enter "HELP FILE NAMES" at an arrow prompt (=>) for a list of files that are available. If you have requested multiple files, you can specify a corrected file name or you can enter "IGNORE" to continue accessing the remaining file names entered.

ENTER A FILE NAME OR (IGNORE): ignore

COST IN U.S. DOLLARS

SINCE FILE TOTAL

ENTRY SESSION

FULL ESTIMATED COST

0.21 0.21

FILE 'AGRICOLA' ENTERED AT 17:03:22 ON 25 MAY 2004

FILE 'BIOTECHNO' ENTERED AT 17:03:22 ON 25 MAY 2004 COPYRIGHT (C) 2004 Elsevier Science B.V., Amsterdam. All rights reserved.

FILE 'CONFSCI' ENTERED AT 17:03:22 ON 25 MAY 2004 COPYRIGHT (C) 2004 Cambridge Scientific Abstracts (CSA)

FILE 'HEALSAFE' ENTERED AT 17:03:22 ON 25 MAY 2004 COPYRIGHT (C) 2004 Cambridge Scientific Abstracts (CSA)

FILE 'IMSDRUGCONF' ENTERED AT 17:03:22 ON 25 MAY 2004 COPYRIGHT (C) 2004 IMSWORLD Publications Ltd.

FILE 'LIFESCI' ENTERED AT 17:03:22 ON 25 MAY 2004 COPYRIGHT (C) 2004 Cambridge Scientific Abstracts (CSA)

FILE 'MEDICONF' ENTERED AT 17:03:22 ON 25 MAY 2004 COPYRIGHT (c) 2004 FAIRBASE Datenbank GmbH, Hannover, Germany

FILE 'PASCAL' ENTERED AT 17:03:22 ON 25 MAY 2004
Any reproduction or dissemination in part or in full,
by means of any process and on any support whatsoever
is prohibited without the prior written agreement of INIST-CNRS.
COPYRIGHT (C) 2004 INIST-CNRS. All rights reserved.

=> (Rh blood)(P)(immobilized or immobilize or immobilization or chromatography)

L1 0 FILE AGRICOLA

PROXIMITY OPERATOR LEVEL NOT CONSISTENT WITH

FIELD CODE - 'AND' OPERATOR ASSUMED 'BLOOD) (P) (IMMOBILIZ'

L2 1 FILE BIOTECHNO L3 0 FILE CONFSCI

L4 0 FILE HEALSAFE

L5 0 FILE IMSDRUGCONF

1 FILE LIFESCI

PROXIMITY OPERATOR LEVEL NOT CONSISTENT WITH

FIELD CODE - 'AND' OPERATOR ASSUMED 'BLOOD) (P) (IMMOBILIZ'

L7 0 FILE MEDICONF

PROXIMITY OPERATOR LEVEL NOT CONSISTENT WITH

FIELD CODE - 'AND' OPERATOR ASSUMED 'BLOOD) (P) (IMMOBILIZ'

L8 0 FILE PASCAL

TOTAL FOR ALL FILES

L9 2 (RH BLOOD) (P) (IMMOBILIZED OR IMMOBILIZE OR IMMOBILIZATION OR CHROMATOGRAPHY)

=> d 19 ibib abs total

L9 ANSWER 1 OF 2 BIOTECHNO COPYRIGHT 2004 Elsevier Science B.V. on STN

ACCESSION NUMBER: 2003:36522970 BIOTECHNO

TITLE: Establishment of human heterohybridoma and

lymphoblastoid cell lines specific for the Rh D and C

antigens

AUTHOR: Pasha R.P.K.; Roohit A.; Shokri F.

CORPORATE SOURCE: Dr. F. Shokri, Department of Immunology, School of

Public Health, Tehran Univ. of Medical Sciences,

Tehran 6446-14155, Iran. E-mail: fazshok@yahoo.com

SOURCE: Transfusion Medicine, (2003), 13/2 (83-92), 34

reference(s)

CODEN: TRMDET ISSN: 0958-7578

DOCUMENT TYPE: Journal; Article COUNTRY: United Kingdom

LANGUAGE: English
SUMMARY LANGUAGE: English
AN 2003:36522970 BIOTECHNO

Human monoclonal antibodies specific for the D antigen of the Rh system AB are valuable tools for blood group typing and prevention of erythroblastosis. In this study, peripheral blood lymphocytes obtained from an Rh-negative woman immunized with Rh-positive fetuses were immortalized with Epstein-Barr virus (EBV), and transformed lymphoblastoid cell lines (LCLs) secreting antibodies to Rh antigens were generated. The presence of specific antibody was assessed by direct haemagglutination using Rh-positive, papain-treated red blood cells (RBCs), and the production of human antibody was assayed by enzyme-linked immunosorbent assay (ELISA). Specificities of the antibodies were determined by a panel of RBCs of known Rh phenotypes. Five LCLs produced antibody specific for the D antigen, and one LCL showed specificity towards the C antigen of the Rh blood group system. High-titre anti-Rh antibody-producing LCLs were subsequently selected and fused with a human x mouse heteromyeloma cell line. A hybridoma line producing human antibody of the immunoglobulin M (IgM) isotype, which strongly reacted with the D antigen, was established. The hybridoma was cloned, and the monoclone has been stable for growth and antibody production during 8 months of continuous culture, with a mean antibody concentration of 11.5 µgmL.sup.-.sup.1 and haemagglutination titre of 1/20 480. This antibody was not able to agglutinate a sample of native

anti-body, followed by Western blot analysis, did not reveal any immobilized D-specific polypeptide. As this human antibody readily agglutinates D.sup.+ RBCs in saline, it has the potential to be used as an efficient reagent in routine blood group typing.

weak D RBCs (D.sup.u); however, agglutination was achieved with papain-treated Du RBCs. Immunoprecipitation of the D antigen by this

L9 ANSWER 2 OF 2 LIFESCI COPYRIGHT 2004 CSA on STN

ACCESSION NUMBER: 88:23744 LIFESCI

TITLE: Characterization of the D, c, E and G antigens of the Rh

blood group system with human monoclonal antibodies.

AUTHOR: Bloy, C.; Blanchard, D.; Lambin, P.; Goossens, D.; Rouger,

P.; Salmon, C.; Masouredis, S.P.; Cartron, J.-P.

CORPORATE SOURCE: Inst. Natl. Transfusion Sanguine and Unite INSERM U76, 6

Rue Alexandre Cabanel, 75015 Paris, France

SOURCE: MOL. IMMUNOL., (1988) vol. 25, no. 9, pp. 925-930.

DOCUMENT TYPE: Journal FILE SEGMENT: F

LANGUAGE: English SUMMARY LANGUAGE: English

The human MAbs, anti-D, -c, -E and -G of the Rh blood group system, produced by Epstein-Barr virus transformed B-cell lines, were purified by protein A-Sepharose chromatography and used to characterize the Rh antigens of human red cells. Scatchard plot analyses performed with the radiolabelled MABs indicated that each R sub(2)R sub(2) red cell carries 0.43, 0.32 and 0.38 x 10 super(5)D, c and E binding sites, respectively. The immune complexes involving anti-c, -E or -G antibodies could be formed with the detergent lysates from red cell

membranes. In contrast, membrane integrity was a prerequisite for the binding of the anti-D antibodies. Finally, from extraction studies of immunocomplexes with non-ionic detergents it was concluded that all the Rh-active components are bound to the membrane skeleton, suggesting that these molecules may have important function for maintaining red cell shape and viability.

L Number	Hits	Search Text	DB	Time stamp
1	1	("20020127236").PN.	USPAT;	2004/05/25 16:44
			US-PGPUB;	
			EPO	
2	1	("6551830").PN.	USPAT;	2004/05/25 16:48
			US-PGPUB;	
			EPO	
3	531	Rh samé chromatography	USPAT;	2004/05/25 16:49
			EPO;	
			DERWENT	
4	7	(Rh near5 blood) same chromatography	USPAT;	2004/05/25 16:51
			EPO;	
			DERWENT	
5	0	(Rh near5 blood) near15 immoboli\$5	USPAT;	2004/05/25 16:51
_	_	,,	EPO;	
			DERWENT	
6	5	(Rh near5 blood) near15 immobili\$5	USPAT;	2004/05/25 16:54
		(III nears proces, nearro manostra,	EPO;	
			DERWENT	
7	5	(Rh near5 blood) same immobili\$5	USPAT:	2004/05/25 16:54
		(in nears brook) bame ramobring	EPO;	2001, 00, 20 10101
			DERWENT	
8	7	(Rh near5 blood) same immobili\$5	USPAT;	2004/05/25 16:56
O	·	(Mi nears brood) same immosiriys	US-PGPUB;	2001,03,23 10.30
			EPO;	
			DERWENT	
9	1 1	("20020127236").PN.	USPAT;	2004/05/25 16:59
9	1	( 2002012/230 ).IN.	US-PGPUB;	2004703723 10.33
	1		EPO EPO	
10	1	("5087338").PN.	USPAT;	2004/05/25 17:00
TO	1	( 3007330 ).EN.	US-PGPUB;	2004/03/23 17:00
			EPO	
1 1	0	(Dh nearE antiqué2 near7 blood) some	USPAT;	2004/05/25 17:02
11	l o	(IIII IIIII)	EPO;	2004/03/25 17:02
		chromat\$5		
10	_	(7)	DERWENT	2004/05/25 17-02
12	6	\	USPAT;	2004/05/25 17:02
		chromatography	EPO;	Į l
4.0	_	(2) 5 11 22	DERWENT	2004/05/25 13 00
13	5	(Rh near5 antige\$3 near7 blood) same	USPAT;	2004/05/25 17:02
		immobili\$3	EPO;	
	İ		DERWENT	1